

Select alternative views for the selected studies

Back to the main page

DISPLAY FORMAT:

- Article list
- Article list
- Text summary**
- Comparison table
- Efficacy graph
- Side effects graph

(MAIN MENU)

SELECT	TITLE	POPULATION	STUDY_ARMS
<input type="checkbox"/>	Efficacy and safety of vildagliptin in patients with type 2 diabetes mellitus inadequately controlled with dual combination of metformin and sulphonylurea.	Type 2 diabetes mellitus (DM2) on Metformin (1500 mg/day or more) plus Glimepiride (4mg/day or more)	<ol style="list-style-type: none"> <li>1. Placebo</li> <li>2. Vildagliptin 50 mg Twice Daily + Metformin 1500 mg/day or more + Glimepiride 4 mg/day or more</li> </ol>
<input type="checkbox"/>	Dapagliflozin, Metformin XR, or both: initial pharmacotherapy for Type 2 Diabetes, a randomised control trial ( <b>*This study includes two trials</b> )	DM2 with hemoglobin A1c (HbA1c) 7.5-12%	<p><b>Trial One:</b></p> <ol style="list-style-type: none"> <li>1. Metformin 500mg Twice Daily + Placebo</li> <li>2. Dapagliflozin 5mg Daily + Placebo</li> <li>3. Dapagliflozin 5mg Daily + Metformin XR</li> </ol> <p><b>Trial Two:</b></p> <ol style="list-style-type: none"> <li>1. Metformin 500mg Twice Daily + Placebo</li> <li>2. Dapagliflozin 10mg Daily + Placebo</li> <li>3. Dapagliflozin 10mg Daily + Metformin XR</li> </ol>
<input type="checkbox"/>	Twice-daily dapagliflozin co-administered with metformin in type 2 diabetes: a 16-week randomized, placebo-controlled clinical trial	DM2 on Metformin (1500mg/day or more) HbA1c: 6.7 - 10.5%	<ol style="list-style-type: none"> <li>1. Placebo + Metformin</li> <li>2. Dapagliflozin 10 mg Daily + Metformin</li> <li>3. Dapagliflozin 5 mg Twice Daily + Metformin</li> <li>4. Dapagliflozin 2.5 mg Twice Daily + Metformin</li> </ol>
<input type="checkbox"/>	Efficacy and safety of liraglutide versus sitagliptin, both in combination with metformin, in Chinese patients with type 2 diabetes: a 26-week, open-label, randomised, active comparator clinical trial	DM2 on Metformin (1000mg/day or more) HbA1c: 7.0-10.0 %	<ol style="list-style-type: none"> <li>1. Liraglutide (inj) 1.8 mg Daily + Metformin</li> <li>2. Sitagliptin 100mg Daily + Metformin</li> </ol>
<input type="checkbox"/>	Efficacy of empagliflozin, sitagliptin, and metformin: an active-comparator, randomized, 78-week open-label extension study in patients with type 2 diabetes.	DM2 who completed one of 12 week double blind randomized controlled trial (DBRCT) HbA1c: 7-10 %	<ol style="list-style-type: none"> <li>1. Metformin</li> <li>2. Empagliflozin 10 mg + Metformin</li> <li>3. Empagliflozin 25 mg + Metformin</li> <li>4. Sitagliptin + Metformin</li> <li>5. Empagliflozin 10 mg + Metformin</li> </ol>

Select studies for comparison